

Determination of the bi-substrate kinetic coefficients for the β -D-glucose- NAD-GDH enzymatic reaction

J. Berthier¹, M. Cubizolles¹, P. Laurent¹, V. Poher², P. Pouteau², M-L. Cosnier¹

¹University Grenoble Alpes, F-38000, Grenoble,

Department of Biotechnology, CEA, LETI, MINATEC, 38054 Grenoble, France

²Avalun, Grenoble, France

ABSTRACT

Colorimetric detection of glucose in sample liquids such as human plasma is made by using enzymatic reactions. In the multi reactional scheme, the first enzymatic reaction is determinant. Either glucose oxidase (GOX) or glucose dehydrogenase (GDH) can be used to convert glucose. We focused here on the study of the enzyme GDH together with the enzymatic cofactor NAD (nicotinamide adenine dinucleotide, oxidized form).


This reaction falls in the category of ternary enzymatic reactions. Such reactions depend on four parameters. A method to determine these four parameters is presented in this work, based on a comparison between a series of experiments and the theory. The best values of the parameters are indicated.

Keywords: ternary enzymatic reaction, bi-substrate kinetics, glucose.

INTRODUCTION

Many glucose tests are now currently available on the market. However, the accuracy of detection can still be improved. The most convenient detection is colorimetry. Colorimetric detection of glucose in sample liquids such as human plasma is made by using enzymatic reactions. In the multi reactional scheme, the first enzymatic reaction is determinant. Either glucose oxidase (GOX) or glucose dehydrogenase (GDH)

can be used to convert glucose. We focused here on the study of the enzyme GDH together with the enzymatic cofactor NAD (nicotinamide adenine dinucleotide). The reaction can be written as



This reaction falls in the category of ternary enzymatic reactions (figure 1). In a first reactional step, the coenzyme NAD (nicotinamide adenine dinucleotide) binds to the GDH (glutamate dehydrogenase) and in a second step, the β -glucose binds to the enzymatic complex. The reaction then occurs, and the product (glucono-lactone) is released together with the NADH (nicotinamide adenine dinucleotide dehydrogenase, reduced form).




Fig.1. Principle of enzymatic ternary reaction: A= β -glucose, B=NAD, P= glucono-lactone, Q= NADH.

Ternary reaction models—also called bi-substrate models, because one enzyme and one coenzyme intervene—depend on four parameters [1-4].

In order to optimize the reaction, these parameters must be known. A method to determine these four parameters is presented in this work. The determination of the four reaction parameters stems from a comparison between a series of experiments and the theory. The best values of the parameters are indicated.

THEORY

The theory of ternary enzymatic reaction is well documented. It is derived from the Michaelis-Menten approach [1-4]. The reaction velocity is given by

$$V = \frac{K_0 [E_0]}{1 + \frac{K_{MB}}{[B]}} \frac{[A]}{[A] + \frac{K_{MA}[B] + K_{SA}K_{MB}}{[B] + K_{MB}}} \quad (2)$$

where $[E]$ is the concentration of enzyme (GDH), $[E_0]$ is the initial enzyme concentration ($[E_0]=1.4 \cdot 10^{-5}$ mM); $[A]$ is the β -D-glucose concentration, $[B]$ the NAD concentration, and the coefficients K 's are the Michaelis-Menten constants. In our case the NAD concentration is fixed and equal to $[B_0] = 0.4$ mM.

Relation (2) generalizes the Michaelis-Menten law: if we note

$$V_{\max,app} = (K_0 [E_0]) / \left(1 + \frac{K_{MB}}{[B]}\right) \quad (3)$$

and

$$K_{M,app} = \frac{K_{MA}[B] + K_{SA}K_{MB}}{([B] + K_{MB})}, \quad (4)$$

relation (2) reduces to the Michaelis-Menten law.

For the concentration $[B]$ the reaction can be simply characterized by the two Michaelis-Menten constants V_{\max} and K_M . But the knowledge of the four ternary reaction constant bares a larger generality.

Reaction kinetics (2) is valid if both concentrations $[A]$ and $[B]$ are such that $[A] \gg [E_0]$ and $[B] \gg [E_0]$, which is our case for reaction (1) and which has been reproduced in the experimental set-up.

MODEL

Relation (2) has three unknowns V , $[A]$ and $[B]$, and four parameters K_0 , K_{MA} , K_{MB} and K_{SA} . By definition the reaction velocity is

$$V = \frac{d[A]}{dt} \quad (5)$$

Using the law of mass action [1,2], we can write

$$[B] = [B_0] - [A_0] + [A] \quad (6)$$

Relation (2) can then be cast in the form of a differential equation in $[A]$, with the initial values $[E_0]$, $[B_0]$

$$\frac{d[A]}{dt} = \frac{K_0 [E_0] \{ [A]^2 + [A]([B_0] - [A_0]) \}}{[A]^2 + [A]([B_0] - [A_0] + K_{MA} + K_{MB}) + K_{SA}K_{MB} + K_{MA}([B_0] - [A_0])} \quad (7)$$

Equation (7) is of the form:

$$\frac{d[A]}{dt} = -cste \frac{P_2([A])}{Q_2([A])}, \quad (8)$$

where P_2 and Q_2 are two second order polynomials.

We have programmed the integration of (8) with the software MATLAB, and used a conventional Runge-Kutta approach to solve for the NAD concentration.

Finally, the product (glucono-lactone) concentration $[Q]$ is given by

$$[Q] = [A_0] - [A] \quad (9)$$

EXPERIMENTS

A series of six kinetics using increasing concentrations of β -D-glucose has been performed (figure 2). Briefly, final concentrations of β -D-glucose, ranging from 0mM to 20mM, were incubated together with NAD and glucose dehydrogenase (*Pseudomonas sp* from Sigma Aldrich) at concentrations of 0.4 mM and 0.5U/mL respectively, using a 25mM Tris, 192mM Glycine buffer pH 8.05 at 37°C. For each β -D-glucose concentration, the reactions have

been done in triplicates, Kinetics were obtained by following the OD (optical density) at 340nm, corresponding to NADH spectrum.




Fig.2. Reaction kinetics for different concentration of β -D-glucose {2,3,6,10,15,20 mM} in Tris/Glycine buffer.

In a Michaelis-Menten type of reaction, the reaction is characterized by the affinity of the substrate (β -D-glucose) for the enzyme (k_M) and the enzyme turn over (k_{cat}), which represent a global view of the reaction.

On the basis of the Lineweaver-Burk plot (figure 3), the affinity of the β -D-glucose for the enzyme can be determined: $k_M = 9.0$ mM. In addition, the enzyme turn over has the value $k_{cat} = 89$ s⁻¹.




Fig.3. Lineweaver-Burke plot used for the determination of k_M and k_{cat} . The slope of the linear fit is k_M/V_{max} , and the intercept with the vertical axis is $1/V_{max}$. The notation [S] stands for the β -D-glucose concentration.

The ratio k_{cat} / k_M , reflecting enzyme efficiency, equals to 9.8×10^3 M⁻¹.s⁻¹. These results are in the same order of magnitude with that obtained in previous works at pH 8.0, performed on wild type

and mutants of GDH isolated from *Bacillus megaterium*, known to be specific for glucose determination in body fluids [5]. In these studies, k_M values ranged from 2.7 to 55 mM and k_{cat} values are comprised between 23 and 430 s⁻¹ [6-8]. Depending on the mutants, the enzyme efficiency previously reported varies from 3.6×10^3 to 42.8×10^3 M⁻¹.s⁻¹. The results that we obtain for the enzyme efficiency using GDH from *Pseudomonas sp* compare well with the previous ones reported on the various forms of GDH from *Bacillus megaterium*.

DETERMINATION OF THE REACTION PARAMETERS

In reality, the reaction is a Michaelis-Menten ternary reaction. In order to characterize the reaction in details, we follow a double reciprocal plot approach. The velocities V are calculated from the kinetic curves of figure 2. In a $\{1/[A], 1/V\}$ coordinates plot, the experimental data points are satisfactorily aligned (figure 4).




Fig.4. Representation of the experimental velocities in a double reciprocal diagram.

The four bi-substrate Michaelis-Menten parameters (K_0 , K_{MA} , K_{MB} and K_{SA}) can be determined by comparison with experimental results.

A linear fit with the experimental results produces the values of the four reaction constants (figure 5). The fitted values of the four bi-substrate reaction constants are: $K_0 = 1.1 \times 10^5$ s⁻¹, $K_{MA} = 1$ mM, $K_{MB} = 2.9$ mM, $K_{SA} = 12$ mM. Reporting these values in (3) and (4), with

$[B]=0.4$ mM, we find $K_{M,app}= 10.7$ mM, and $V_{max,app}= 0.003$ mM/s, and $k_{cat,app}= V_{max,app} / [E_0]= 222$ s⁻¹. The values of apparent K_M and k_{cat} deduced from the ternary model are in the same range as the corresponding experimental Michaelis-Menten kinetic constants.




Fig.5. Fit of the theoretical model on the experimental results in the double Lineweaver-Burke diagram.

Reversing the approach, and reporting the fitted parameters in relation (5), we obtain a satisfactory comparison of the kinetics between experiments and model (figure 6), confirming the approach.




Fig.6. Kinetics comparison between model and experiments: continuous lines correspond to the experimental measures and the dots to the model.

CONCLUSION

In this work, we propose a method for the determination of the kinetic coefficients for the β -D-glucose-NAD-GDH enzymatic ternary reaction, based on the comparison between the theory of bi-substrate enzymatic reaction and experiments.

In order to determine the four constants of the reaction, we place ourselves in a double reciprocal plot approach $\{1/[A], 1/V\}$ where the kinetic curves are linear. Such an approach facilitates the fit of the constants. A reverse reconstruction reproduces the kinetic signals.

The knowledge of the coefficients of the reaction is essential to optimize glucose detection systems. Moreover, the method can be generalized to other bi-substrate reactions.

REFERENCES

- [1] I.H. Segal, Enzyme kinetics, John Wiley & Sons, New York, 1975.
- [2] A. Cornish-Bowden, Fundamentals of enzyme kinetics, 3rd Edition, 2004, Portland Press
- [3] P.A. Frey, A.D. Hegeman Enzymatic reaction mechanisms, Oxford University Press, New York, 2007.
- [4] C.A. Janson, W.W. Cleland, The kinetic mechanism of glycerokinase, J Biol Chem. **249**(8), 2562-2566, 1974.
- [5] H. Rindfrey, R. Helger and H. Lang, Kinetic Determination of Glucose Concentrations with Glucose Dehydrogenase, J. Clin. Chem. Clin. Biochem., **15**, 217-220, 1977.
- [6] S.-H. Baik, T. Ide, H. Yoshida, O. Kagami, S. Harayama, Significantly enhanced stability of glucose dehydrogenase by directed evolution, Appl Microbiol Biotechnol **61**:329–335, 2003.
- [7] T. Nagao, T. Mitamura, Xiao Hui Wang, S. Negoro, T. Yomo, I. Urabe, H. Okada, Cloning, Nucleotide Sequences, and Enzymatic Properties of Glucose Dehydrogenase Isozymes from Bacillus megaterium IAM1030, J. Bacteriology, **174**(15), 5013-5020, 1992.
- [8] K. Yamamoto, T. Nagao, Y. Makino, I. Urabe, H. Okada, Characterization of Mutant Glucose Dehydrogenases with Increasing Stability, Ann N Y Acad Sci. **613**, 362-365, 1990.